



Epidemiological Profile of Substance Use + Related Factors in Minnesota's American Indian + Alaska Native Communities

Developed by EpiMachine, LLC on behalf of the Minnesota State Epidemiological
Outcomes Workgroup

April 2017

TABLE OF CONTENTS

| | |
|---|-----------|
| INTRODUCTION | 3 |
| POPULATION SNAPSHOT | 5 |
| KEY FINDINGS | 6 |
| ALCOHOL | 7 |
| TOBACCO | 11 |
| DRUGS | 13 |
| MENTAL HEALTH | 18 |
| SHARED RISK + PROTECTIVE FACTORS | 20 |
| DATA SOURCES | 24 |

INTRODUCTION

MINNESOTA STATE EPIDEMIOLOGICAL OUTCOMES WORKGROUP

The Minnesota State Epidemiological Outcomes Workgroup (SEOW) has been monitoring trends in substance use and related problems since 2007. The SEOW is funded by the Substance Abuse and Mental Health Services Administration (SAMHSA) Center for Substance Abuse Prevention (CSAP), led by the Minnesota Department of Human Services (DHS) Alcohol and Drug Abuse Division (ADAD), and staff through a subcontract with EpiMachine, LLC.

SEOW membership includes representation from the Minnesota Department of Human Services, Minnesota Department of Health, Minnesota Department of Education, Minnesota Board of Pharmacy, Wilder Research, the Hazelden Betty Ford Graduate School of Addiction Studies, and EpiMachine. Project staff manage and maintain the online substance abuse prevention data query system SUMN.org, provide training and technical assistance to communities across Minnesota, and develop a variety of data products—including this community profile.

SUMN.org

The Substance Use in Minnesota or SUMN.org website houses county, regional, and state data from a number of sources on:

- Alcohol use, consequences, and related factors
- Tobacco use, consequences, and related factors
- Drug use, consequences, and related factors
- Mental health
- Risk and protective factors shared between substance use and mental health

Site visitors can produce tables, maps, graphs, and charts using Data by Topic. Visitors can also search available data by location, or by demographic group. In addition to key prevention data, SUMN.org also features Publications, Community Resources, Toolbox, and Gallery. The Publications page includes SEOW data products, and links to state and national reports, articles, and websites. The Community Resources page provides links and contact information on coalitions and prevention organizations across Minnesota. The Toolbox includes tip sheets, guidance documents, training materials, and toolkits related to data collection, analysis, translation, formatting, dissemination, and use. The Gallery features examples of fact sheets, posters, public service announcements, infographics, and more created by visitors using SUMN data.

COMMUNITY PROFILES

This profile is grounded in CSAP's Strategic Prevention Framework (SPF). The SPF is a five-step prevention planning model consisting of 1) Assessment (of both need and resources), 2) Capacity Building, 3) Planning, 4) Implementation, and 5) Evaluation. The profile serves as an integral step in the assessment phase of the SPF. It has been created to summarize substance use among American Indians and Alaska Natives in Minnesota, and related factors.

This profile was created to help the state and communities determine prevention needs based upon available data on substance use and related factors. Accordingly, the profile can be used by a variety of audiences for related, but different, purposes. State-level administrators may use the profile to prepare applications for federal funding or to monitor prevention-related trends in local communities to which they administer grants. Community-level prevention planners may use it to assess the relative importance of substance related problems in their communities or to apply for grant funding themselves. Overall, the profile is intended to help all audiences in Minnesota make decisions based on existing evidence and demonstration of need.

It is important for state, county and city planners to have accurate and readily available data on American Indian and Alaska Native substance use and consequences—and for all communities—in order to paint a complete picture of need in our state. Aggregated data do not reveal disparities that exist in a given location or for a specific population. While overall use of a substance may be low in Minnesota, it could be quite high within a particular community or population. Community-specific data allows for well-planned and targeted interventions. Every effort should be made at the national-, state-, county- and city-level to collect data by race/ethnicity. It is also important to recognize the limits of broad race and ethnicity categories.

This profile can be used by community leaders and prevention professionals to plan, set priorities, target resources, and simply to spur conversation about community-level alcohol, tobacco and other drug use and consequences. The goal of this profile, and the State Epi Profile, is to encourage data-driven decision making over reliance on anecdotal information. This report is by no means exhaustive. Community leaders and prevention professionals can use this profile in conjunction with community-level data and qualitative information from surveys, focus groups and key informant interviews.

There are some important limitations and data gaps to note. Race selections on surveys do not always allow for self-report of specific tribal affiliation or membership. Racial designations made on death and arrest reports are often done by medical examiners and law enforcement, and therefore may not be accurate. Race/ethnicity is not always labeled or defined uniformly; some sources report for —mixed race while others report for American Indian and Alaska Native alone or in combination with one or more races/ethnicities. County and city level data on American Indians and Alaska Natives are often not available due to small numbers and/or low survey response rate, if collected at all. Also, rates may fluctuate greatly due to small numbers.

POPULATION SNAPSHOT

According to the 2015 American Community Survey, 2% of Minnesotans identify as American Indian and Alaska Native alone (alone or in combination with other races/ethnicities). American Indian and Alaska Native communities in Minnesota are considerably younger than the state average, with over one-third being under the age of 18.

| 2015 | American Indians +Alaska Natives in Minnesota | | All Minnesotans | |
|-----------------|---|---------|-----------------|---------|
| | Number | Percent | Number | Percent |
| Under 18 years | 42,655 | 36.3% | 1,360,354 | 24.2% |
| 18 to 24 years | 14,250 | 12.1% | 525,404 | 9.3% |
| 25 to 44 years | 30,867 | 26.3% | 1,450,416 | 25.8% |
| 45 to 64 years | 22,478 | 19.1% | 1,485,654 | 26.4% |
| 65 years + over | 7,314 | 6.2% | 809,397 | 14.4% |
| Total | 117,564 | | 5,631,225 | |

Annual Estimates of the Resident Population by Sex, Age, Race Alone or in Combination, and Hispanic Origin for the United States and States: April 1, 2010 to July 1, 2015 
2015 Population Estimates

There are two tribes located in Minnesota, the Sioux and Ojibwe; four nations in the Sioux tribe and seven nations in the Ojibwe tribe. Members of other tribes have moved to Minnesota as well.

According to the Minnesota State Demographic Center's Young Adults in Minnesota: A Demographic & Economic Profile, 1 of every 100 young adults (ages 18-34) is American Indian. https://mn.gov/admin/assets/young-adults-in-mn-profile-popnotes-june2015_tcm36-219658.pdf

The Minnesota Student Survey provides data for a number of indicators in this profile. In 2016, the overall survey sample identifying American Indian was 2,622.

| Number of 2016 Minnesota Student Survey Respondents Identifying as non-Hispanic American Indian | | | | |
|--|-------|--------|-----------|-------|
| | Male | Female | No answer | Total |
| 5 th Grade | 633 | 467 | 6 | 1,106 |
| 8 th Grade | 383 | 273 | 0 | 656 |
| 9 th Grade | 322 | 215 | 2 | 539 |
| 11 th Grade | 177 | 140 | 4 | 321 |
| Total | 1,515 | 1,095 | 12 | 2,622 |

KEY FINDINGS

POSITIVE FINDINGS

- The number of American Indians and Alaska Natives arrested for driving under the influence or for liquor laws has been decreasing over time (MN BCA)
- The percent of American Indian/Alaska Native students reporting past 30 day alcohol use and binge drinking decreased from 2013 to 2016 (MSS)
- While still higher than average, American Indian and Alaska Native adult rates of binge drinking, cigarette use, and depressive symptoms are down from 2010 (2015 MNSASU)

AREAS FOR IMPROVEMENT


- American Indian/Alaska Native students were less likely than the state average to perceive great or moderate risk of harm from using alcohol, cigarettes, marijuana, or prescription drugs (2016 MSS)
- American Indian/Alaska Native students were more likely than the state average to report past 30 day alcohol use, 19% vs. 14%, binge drinking, 11% vs. 6%, tobacco/nicotine use, 23% vs. 13%, marijuana use, 25% vs. 8%, and prescription drug misuse: 9% vs. 5% (2016 MSS)
- American Indian/Alaska Native adults were more likely than average to report past 30 day binge drinking, cigarette and e-cigarette smoking, marijuana use, and past two week depressive symptoms; they are also less likely to perceive great or moderate risk of harm from substance use (2015 MNSASU)
- The percentage of American Indians and Alaska Natives admitted to Minnesota treatment programs for opioids as their primary substance of abuse has risen sharply, as have admissions for methamphetamines (DAANES)
- American Indian/Alaska Native students were more likely than the state average to report past year suicidal ideation, 19% vs. 12%, and past year suicide attempts: 9% vs. 4% (2016 MSS)
- American Indian/Alaska Native students were more likely than the state average to report high distress levels for both internalizing and externalizing disorders (2013 MSS)
- American Indian/Alaska Native students were more likely than the state average to report one or more adverse childhood experiences (2016 MSS)
- American Indian/Alaska Native students were less likely than the state average to feel safe at school, feel safe going to and from school, or feel safe in their neighborhood (2016 MSS)
- American Indian/Alaska Native students were less likely than the state average to feel that their parents, other adults, friends, teachers and other adults at school, and adults in their community care about them “quite a bit” or “very much” (2016 MSS)

ALCOHOL

YOUTH USE

Current alcohol use is measured by the Minnesota Student Survey as students reporting any use of alcohol in the past 30 days. In 2016, 18.8% of American Indian and Alaska Native (non-Hispanic) 8th, 9th, and 11th grade students reported current alcohol use (down from 24.2% in 2013) as compared to the state average of 13.9% (down from 16.8% in 2013).


American Indian and Alaska Native 8th, 9th, and 11th grade females were more likely than males to report current alcohol use in 2016: 22.1% vs. 16.4%. American Indian and Alaska Native 5th graders were a bit more likely than the state average to report any use of alcohol in the last 12 months: 4.6% vs. 3.4%. American Indian and Alaska Native students were more likely than the state average to report past 30 day alcohol use regardless of grade level (see graph below).



American Indian and Alaska Native 8th, 9th, and 11th grade students living outside the seven-county metro area were more likely to report current alcohol use as compared to those living in the metro: 20.4% vs. 14.6%.

Binge drinking is measured by the Minnesota Student Survey as students reporting having five or more drinks in a row on one occasion within the past 30 days. In 2016, 11.2% of American Indian and Alaska Native (non-Hispanic) 8th, 9th, and 11th grade students reported binge drinking (down from 14.9% in 2013) as compared to the state average of 6.2% (down from 8.2% in 2013).

American Indian and Alaska Native 8th, 9th, and 11th grade females were more likely to report binge drinking in 2016 as compared to males: 12.8% vs. 10.0%. American Indian and Alaska Native students were more likely than the state average to report binge drinking, regardless of grade level (see graph at top of next page).



American Indian and Alaska Native 8th, 9th, and 11th grade students living outside the seven-county metro area were more likely to report binge drinking as compared to those living in the metro: 12.5% vs. 7.6%.


ADULT USE

Binge drinking is measured by the Minnesota Survey of Adult Substance Use as having five or more drinks (four or more for women) in a row on one occasion within the past 30 days. In 2015, 16.1% of American Indian and Alaska Native adults in Minnesota reported binge drinking (down from 20.3% in 2010) as compared to the state average of 13.9% (down from 18.2% in 2010).

CONSEQUENCES

Data from the Centers for Disease Control and Prevention Compressed Mortality File show that age-adjusted alcohol-related deaths among American Indians and Alaska Natives in Minnesota have been consistently higher than the state average. The average age-adjusted alcohol-related death rate per 100,000 population from 2006 to 2015 was 44.2 for American Indians and Alaska Natives and 7.9 for the state as a whole. Alcohol-induced causes of death were determined by the National Center for Health Statistics based on analysis by the Selected Cause of Death groups.


The Minnesota Bureau of Criminal Apprehension (BCA) provides data on arrests for driving under the influence (DUI). The BCA also provides data on liquor law arrests. This indicator excludes drunkenness, driving under the influence, and federal violations. It includes state or local violations related to furnishing liquor to a minor, using a vehicle for illegal transportation of liquor, bootlegging, and operating a still. The number of American Indian Minnesotans arrested for liquor laws in 2015 was 494, down from 1,376 in 2006. The number arrested for DUI in 2015 was 309, down from 769 in 2006 (see graph at top of next page).



According to the 2015 American Community Survey estimates, 2% of Minnesotans identify as American Indian/Alaska Native. In 2015, 5% of Minnesotans arrested for liquor laws were AI/AN, and 1.5% arrested for DUI were AI/AN.

TREATMENT

In 2015, 30.4% of American Indians and Alaska Natives admitted to Minnesota treatment facilities were admitted for alcohol as their primary substance of abuse (compared to 40.9% for all races) according to the Drug and Alcohol Abuse Normative Evaluation System. The percent of American Indians and Alaska Natives admitted for alcohol, as opposed to other substances, has declined over time as has the state average.




RISK + PROTECTIVE FACTORS

American Indian and Alaska Native 5th, 8th, 9th, and 11th grade students were less likely to report in 2016 that they believe people put themselves at great or moderate risk of harm by frequently binge drinking: 53.1% vs. 71.2%. AI/AN 9th and 11th grade students were considerably more likely than the Minnesota average to report age of first alcohol use at 13 or younger: 28.0% vs. 16.9%.

In 2016, AI/AN 5th, 8th, 9th, and 11th grade students were less likely than average to report their parents would think it's wrong or very wrong for them to have one or two drinks of an alcoholic beverage nearly every day: 84.4% vs. 90.5%. Similarly, they were less likely than average to think their friends would feel it's wrong or very wrong: 74.6% vs. 81.6%.

Youth tend to misperceive the number of peers who are drinking. Those who perceive that most students drink monthly or more often are more likely to report they drank in the past 30 days themselves. Among AI/AN 8th, 9th, and 11th grade students, 31.7% think most students in their school have never had alcohol. However, 70.3% say they themselves have never had alcohol (see graph below).




According to the 2015 Minnesota Survey of Adult Substance, American Indian and Alaska Native adults in Minnesota were less likely than average to believe people put themselves at great risk of harm by binge drinking: 62.6% vs. 69.9%. They were somewhat more likely than average though to believe it's very or somewhat likely that someone drinking and driving would be stopped by the police in their community: 78.8% vs. 75.6%.

TOBACCO

YOUTH USE

In 2016, 22.8% of American Indian and Alaska Native (non-Hispanic) 8th, 9th, and 11th grade students reported any past 30 day tobacco/nicotine use (not including traditional tobacco practices) as compared to the state average of 12.8%. Among the various types of tobacco/nicotine products, AI/AN students were most likely to report use of e-cigarettes followed by standard commercial cigarettes (see graph below).



American Indian and Alaska Native 8th, 9th, and 11th grade females were more likely than males to report any past month tobacco/nicotine use in 2016: 25.1% vs. 21.2%. Among AI/AN students, past 30 day tobacco/nicotine use was reported by 17.2% of 8th graders, 22.3% of 9th graders, and 34.8% of 11th graders.


AI/AN 8th, 9th, and 11th grade students living outside the seven-county metro area were more likely to report any past 30 day tobacco/nicotine use compared to those living in the metro: 24.4% vs. 18.2%.

ADULT USE

According to the 2015 Minnesota Survey of Adult Substance Use, 46.1% of American Indian and Alaska Native adults in Minnesota reported smoking cigarettes on one or more days in the past 30 days (down from 58.9% in 2010) as compared to the state average of 15.7% (down from 19.7% in 2010). American Indian and Alaska Native adults were also considerably more likely than average to report past 30 day use of e-cigarettes: 11.9% vs. 5.3%.

CONSEQUENCES

Data from the Centers for Disease Control and Prevention Compressed Mortality File show that age-adjusted lung, bronchus, and trachea cancer death rates (ICD codes C33 and C34) for American Indians and Alaska Natives in Minnesota has fluctuated over time, but remained consistently higher than the overall state rate. From 2014 to 2015, the rate for AI/AN in Minnesota dipped from 81.8 per 100,000 population to 60.2.



RISK + PROTECTIVE FACTORS


American Indian and Alaska Native 5th, 8th, 9th, and 11th grade students were considerably less likely than the state average to report in 2016 that they believe people put themselves at great or moderate risk of harm by smoking one or more packs of cigarettes per day: 58.3% vs. 78.1%.

According to the 2015 Minnesota Survey of Adult Substance, American Indian and Alaska Native adults in Minnesota were less likely than average to believe people put themselves at moderate or great risk of harm from smoking cigarettes: 78.8% vs. 85.2%. They were also less likely than average to perceive moderate or great risk of harm from using e-cigarettes: 57.3% vs. 60.9%.


DRUGS

YOUTH USE


In 2016, American Indian and Alaska Native (non-Hispanic) 11th graders were more likely than average to report past year alcohol use, marijuana use, and use of all illicit drugs other than marijuana (see graph below).



In 2016, 24.7% of AI/AN (non-Hispanic) 8th, 9th, and 11th grade students reported any past 30 day marijuana as compared to the state average of 8.2%. AI/AN 8th, 9th, and 11th grade females were considerably more likely than males to report current marijuana use in 2016: 31.0% vs. 20.0%. AI/AN students across all grades were more likely than the state average to report current marijuana use (see graph on next page). AI/AN 8th, 9th, and 11th grade students living outside the seven-county metro area were more likely to report current marijuana compared to those living in the metro: 21.5% vs. 12.5%.



In 2016, 9.3% of American Indian and Alaska Native (non-Hispanic) 8th, 9th, and 11th grade students reported any past 30 day use of prescription drugs not prescribed for them (taken only to get high) as compared to the state average of 4.6%. Across all grade levels, AI/AN students were more likely than average to report past month prescription drug misuse (see graph below).



American Indian and Alaska Native 8th, 9th, and 11th grade students were more likely than average to report past 12 month misuse of prescription drugs, regardless of drug type:

- Pain relievers: 5.2% vs. 3.2%
- ADD/ADHD drugs: 4.4% vs. 2.7%
- Tranquilizers: 3.6% vs. 1.8%
- Stimulants: 2.3% vs. 0.9%


ADULT USE

According to the 2015 Minnesota Survey of Adult Substance Use, 9.9% of American Indian and Alaska Native adults in Minnesota reported use of marijuana in the past 30 days (down from 10.5% in 2010) as compared to the state average of 4.8% (up from 4.4% in 2010). In 2010, American Indian and Alaska Native adults in Minnesota were considerably more likely than the state average to report any past 12 month use of illicit drugs other than marijuana: 11.1% vs. 4.6%. While the 2015 survey didn't provide large enough estimates to reliably report illicit drug use by race/ethnicity, the over state-level rate dropped from 4.6% to 1.9%.

CONSEQUENCES

The number of American Indians and Alaska Natives in Minnesota arrested for narcotics has increased over the past five years, from 446 in 2011 to 648 in 2015. About 4% of all Minnesotans arrested for narcotics in 2015 were AI/AN.


Data from the Centers for Disease Control and Prevention Compressed Mortality File show that age-adjusted drug-related deaths (ICD -10 codes X40-44, X60-64, X85 and Y10-14) among American Indians and Alaska Natives in Minnesota have been consistently higher than the state average. The AI/AN drug-related death rate increased sharply from 2010 to 2012, then flattened out. In 2015, age-adjusted drug-related death rate per 100,000 population was 47.3 for AI/ANs and 10.6 for the state as a whole.



TREATMENT

According to the Drug and Alcohol Abuse Normative Evaluation System, American Indians and Alaska Natives admitted to Minnesota treatment programs for a primary substance of abuse other than alcohol were most likely to be admitted for opioids. In prior years marijuana had been the second primary substance of abuse after alcohol. Admissions to treatment for opioids has increased significantly among American Indians and Alaska Natives since 2005, as it has among the overall treatment population (see graph below).


While American Indians and Alaska Natives have been less likely than the state average to be admitted for methamphetamines as their primary substance use abuse, rates of admission are on the rise. In 2015, admissions for American Indians and Alaska Natives were as follows: 15.8% for marijuana, 34.7% for opioids, 16.5% for methamphetamines, and 1.2% for crack/cocaine.



RISK + PROTECTIVE FACTORS

According to the Minnesota Student Survey, American Indian and Alaska Native 5th, 8th, 9th, and 11th grade students were considerably less likely to report in 2016 that they believe people put themselves at great or moderate risk of harm by smoking marijuana once or twice per week as compared to the state average: 42.2% vs. 61.7%. They were also considerably less likely than average to report that they believe people put themselves at great or moderate risk of harm by using prescription drugs not prescribed for them: 60.3% vs. 78.8%.

Youth tend to misperceive the number of peers who are using marijuana. Those who perceive that most students use marijuana monthly or more often are more likely to report they used in the past 30 days themselves. Among AI/AN 8th, 9th, and 11th grade students, 30.7% think most students in their school have never had marijuana. However, 69.9% say they themselves have never had marijuana (see graph below).




According to the 2015 Minnesota Survey of Adult Substance, American Indian and Alaska Native adults in Minnesota were considerably less likely than average to believe people put themselves at moderate or great risk of harm from using marijuana: 46.8% vs. 60.2%. They were also considerably less likely than average to perceive moderate or great risk of harm from misusing prescription drugs: 81.8% vs. 93.9%.

MENTAL HEALTH

We also know from national research literature that substance use is a risk factor for mental health problems, and in turn mental health problems are a risk factor for substance use. We also know that many Minnesotans suffer from co-occurring substance use and mental health disorders.

As an example, Minnesota students who reported seriously considering suicide in the past year, actually attempting suicide in the past year, or harming themselves on purpose in the past year were more likely than those who did not report those experiences to abuse substances in the past 30 days. As shown in the chart below, Minnesota students reporting past year suicidal ideation were over twice as likely to report past 30 day alcohol use and three times more likely to report past 30 day marijuana use.



SUICIDE

Minnesota Student Survey findings from 2016 show that American Indian and Alaska Native 8th, 9th, and 11th grade students were more likely than the state average to report past-year suicidal ideation: 19.1% and 11.8% respectively. They were also more likely to report a past-year suicide attempt: 9.4% and 3.6% respectively.

SELF-HARM

American Indian and Alaska Native 8th, 9th, and 11th grade students were considerably more likely to report any past-year self-harm 25.0% vs. 15.6%. Self-harm includes doing something to purposely injure oneself without want to die, such as cutting, burning, or bruising oneself on purpose.

ADULT MENTAL HEALTH

According to the 2015 Minnesota Survey of Adult Substance Use, American Indian and Alaska Native adults in Minnesota were more likely than the state average to report at least moderate depressive symptoms in the past two weeks: 12.5% vs. 7.4%. Depressive symptoms are based on the 8-item Patient Health Questionnaire (PHQ-8). However, rates for American Indian and Alaska Native adults were down from 21.0% in 2010.


SHARED RISK + PROTECTIVE FACTORS

Risk factors are characteristics at the biological, psychological, family, community, or cultural level that precede and are associated with a higher likelihood of problem outcomes; protective factors are characteristics associated with a lower likelihood of problem outcomes or that reduces the negative impact of a risk factors on problem outcomes. Some risk factors are specifically associated with substance use, such as perceived risk of harm. Some risk and protective factors are associated with both substance use/abuse and with mental health.

ADVERSE CHILDHOOD EXPERIENCES (ACEs)



The Adverse Childhood Experiences (ACE) Study was a collaboration between the Centers for Disease Control and Prevention (CDC) and Kaiser Permanente’s Health Appraisal Clinic in San Diego. The study found that childhood experiences of abuse, neglect, and family dysfunction are linked to leading causes of illness, poor quality life, and death. An “ACE Score” is calculated by adding 1 point for each ACE experienced. An ACE score from the 2016 MSS can range from zero to seven based on the following survey items:

- Parent or guardian who is currently in jail, and/or has been in jail in the past
- Live with someone who drinks too much alcohol
- Live with someone who uses illegal drugs or abuses prescription drugs
- Verbal abuse by a parent or other adult in the household
- Physical abuse by a parent or other adult in the household
- Parents or other adults in the home who physically abuse each other
- Sexual abuse by an older or stronger family member, or someone outside the family



Among American Indian and Alaska Native students responding to the 2016 Minnesota Student Survey, 39.3% reported an ACE score of zero, 29.9% an ACE score of one, 13.5% an ACE score of two, 8.7% an ACE score of three, and 8.6% an ACE score of four or more.

Past 30 day alcohol use increases incrementally with ACE score, as do past year suicidal ideation and attempts. While not shown here, each increase in ACE score is also associated with: increased marijuana use, increased cigarette smoking, lower grades, lower feelings of safety at school, home or in the community, and a decreased perception that parents, friends, teachers, and adults in the community care.



BULLYING

The 2016 Minnesota Student Survey included a number of measures of bullying and harassment. Bullying behaviors experienced at school include:

- Pushed, shoved, slapped, hit or kicked by another student who was not kidding around, *and/or*
- Another student threatened to beat them up, *and/or*
- Spread mean rumors or lies about them, *and/or*
- Made sexual jokes, comments or gestures toward them, *and/or*
- Excluded them from friends, other students, or activities

| | Experienced any past 30 day bullying behavior | Did NOT experience any past 30 day bullying behavior |
|---|---|--|
| Past 30 day alcohol use | 18.4% | 10.2% |
| Past 30 day marijuana use | 11.2% | 6.4% |
| Past 12 month self-harm | 25.3% | 7.8% |
| Past 12 month serious suicide consideration | 19.5% | 5.7% |

Compared to the state average, American Indian and Alaska Native students were a bit more likely to report experiencing bullying in the past 30 days (47.8% vs. 44.2%).

SAFETY


Students who feel safe at school are less likely to report past 30 day substance use, and less likely to report mental health issues. According to the 2016 Minnesota Student Survey, AI/AN students are less likely than the state average to report feeling safe at school—88.5% said they “strongly agree” or “agree” they feel safe at school compared to the state average of 92.7%. They are also less likely to report feeling safe going to and from school: 92.6% vs. 95.6%.

| | Students "strongly agree" or "agree" that they feel safe at school | Students "disagree" or "strongly disagree" that they feel safe at school |
|---|--|--|
| Past 30 day alcohol use | 13.0% | 24.4% |
| Past 30 day marijuana use | 7.9% | 17.1% |
| Past 12 month self-harm | 13.9% | 37.3% |
| Past 12 month serious suicide consideration | 10.2% | 32.0% |

Youth who feel safe in their neighborhoods are also less likely to report past month substance use or past year mental health problems. In 2016, 88.0% of American Indian and Alaska Native students said they “strongly agree” or “agree” that they feel safe in their neighborhood compared to the state average of 95.0%.

SUPPORT

Youth who feel that others care about them are less likely to report substance use and mental health problems. For example, Minnesota students who feel their teachers and other adults at school care about them quite a bit or very much are 2.7 times less likely to report past 30 day alcohol use than those who feel they care about them a little or not all. They are 4.3 times less likely to report suicidal ideation. In 2016, American Indian and Alaska Native students were somewhat less likely than the state average to report feeling that others care about them very much.



DATA SOURCES

MINNESOTA STUDENT SURVEY (MSS)

The MSS is a confidential and anonymous self-administered survey sponsored by the Minnesota schools, the Minnesota Department of Education, the Minnesota Department of Health, the Minnesota Department of Human Services, and the Minnesota Department of Public Safety. The survey is administered every three years to students attending Minnesota public, charter and tribal schools. From 1995 to 2010, the survey was given to 6th, 9th and 12th graders; starting in 2013 the survey was administered to 5th, 8th, 9th, and 11th graders.

<http://www.health.state.mn.us/divs/chs/mss/>

<http://education.state.mn.us/MDE/StuSuc/SafeSch/MNStudentSurvey/>

MINNESOTA SURVEY ON ADULT SUBSTANCE USE (MNSASU)

The MNSASU is a statewide telephone survey conducted by DHS—the last survey was administered in 2014/2015. The primary objective of this project is to obtain current estimates of the number of adults in the general population in Minnesota who are abusing or dependent on alcohol or other drugs and are in need of treatment. The population for this survey included Minnesota residents 18 years of age or older and non-institutionalized.

http://www.dhs.state.mn.us/main/idcplg?IdcService=GET_DYNAMIC_CONVERSION&dID=158864

UNIFORM CRIME REPORTS (UCR)

The Minnesota Bureau of Criminal Apprehension collects activity information from law enforcement agencies throughout the State of Minnesota. Uniform Crime Reports measure the amount of criminal activity within the State as collected and prepared from data submitted by individual law enforcement agencies. The offense categories presented in the Profile are Part II offenses: liquor laws and narcotics arrests. <http://www.dps.state.mn.us/bca/CJIS/Documents>

DRUG AND ALCOHOL ABUSE NORMTATIVE EVALUATION SYSTEM (DAANES)

DAANES includes data on all private- and public-pay treatment facility admissions and discharges. Data were obtained by request from the Performance Measurement and Quality Improvement (PMQI) Division.

CDC COMPRESSED MORTALTY FILE

The Centers for Disease Control and Prevention Compressed Mortality database contains mortality and population counts for all U.S. counties for the years 1979 to 2010. Counts and rates of death can be obtained by underlying cause of death, state, county, age, race, sex, and year. Beginning in 1999, cause of death is specified with the International Classification of Diseases 10th Revision (ICD 10) codes. <http://wonder.cdc.gov/mortSQL.html>