



# FARIBAULT COUNTY

SUMN.org FACT SHEET

2017

## Substance Abuse Prevention

This fact sheet provides county level data on substance use and related factors to help guide decisions about prevention efforts. These data can be used to justify need for prevention funding, monitor substance-related trends, and plan programs and initiatives by establishing community level prevention priorities.


According to 2016 estimates from the U.S. Census Bureau, Faribault County has 13,935 residents; 22% are under the age of 18.

### Key Findings

- Compared to the state average, a larger percentage of Faribault County students reported past 30-day alcohol use, binge drinking, and misuse of prescription drugs (see Fig. 1).
- Cigarette smoking among 9<sup>th</sup> graders has declined from 20% in 2001 to 4% in 2016; however, 7% of 9<sup>th</sup> graders using e-cigarettes in 2016.
- Among Faribault County 8<sup>th</sup>, 9<sup>th</sup>, and 11<sup>th</sup> graders, those with better teacher-student relationships are nearly 2.5 times less likely to report past-month marijuana use.
- Faribault County's alcohol-related motor vehicle crash rate and arrest rate for driving while intoxicated (DWI) are both higher than the state averages.
- From 2000 to 2015, Faribault County recorded 3 overdose deaths due to opioids, and 1 due to heroin.

*For more information on alcohol, tobacco, and other drug use, consequences, contributing factors, and treatment, as well as mental health data, visit [SUMN.org](http://SUMN.org)*

Figure 1. 8<sup>th</sup>, 9<sup>th</sup>, and 11<sup>th</sup> Grade Past 30 Day Substance Use, 2016 Minnesota Student Survey (MSS)



\*\*Tobacco includes any past month use of cigarettes, cigars, chewing tobacco, electronic cigarettes and/or hookah.

## Adult Substance Use

According to the 2015 Minnesota Survey on Adult Substance Use (MNSASU):

- Past-month binge drinking (women having 4+ drinks or men having 5+ drinks in a row on one occasion) was reported by 11% of adults in the Southwest region and 14% of Minnesota adults.
- Past-month cigarette smoking was reported by 16% of Southwest adults and 16% of Minnesota adults.
- Past-month marijuana use was reported by 2% of adults in the Southwest region compared to 5% of adults statewide.

## Youth Substance Use

Data on youth substance use are from the Minnesota Student Survey (MSS); Faribault County students make up 0.3% of the overall statewide sample.

**Table 1. 2016 Minnesota Student Survey Faribault County Respondents**

	5 <sup>th</sup> Grade	8 <sup>th</sup> Grade	9 <sup>th</sup> Grade	11 <sup>th</sup> Grade
Females	51	47	45	55
Males	59	60	63	59
No Answer	0	0	0	0
Total	110	107	108	114

Past year alcohol and/or drug use was reported by 5.1% of Faribault 5<sup>th</sup> graders. Questions about past 30-day substance use (covered in the following section) are not asked of 5<sup>th</sup> grade students.

### Alcohol

Past 30-day alcohol use was reported by 10% of 8<sup>th</sup> graders, 15% of 9<sup>th</sup> graders, and 33% of 11<sup>th</sup> graders in Faribault County in 2016. Use among 9<sup>th</sup> graders dropped in Faribault County from 27% in 2001 to 15% in 2016. Past 30-day binge drinking (defined as having five or more drinks in a row on one occasion) was reported by 1% of 8<sup>th</sup> graders, 4% of 9<sup>th</sup> graders, and 18% of 11<sup>th</sup> graders in 2016.

### Tobacco/Nicotine

Of Faribault County 8<sup>th</sup>, 9<sup>th</sup> and 11<sup>th</sup> graders, 12% reported any past 30-day tobacco use: 6% reported smoking cigarettes, 5% reported smoking cigars or cigarillos, and 9% reported chewing tobacco use. In addition, 9% reported e-cigarette use, and 4% reported hookah use; questions about these two methods of nicotine use were added to the survey in 2016. Cigarette smoking among 9<sup>th</sup> graders dropped in Faribault County from 20% in 2001 to 4% in 2016.

### Marijuana

Past 30-day marijuana use was reported by 2% of 8<sup>th</sup> graders, 7% of 9<sup>th</sup> graders, and 17% of 11<sup>th</sup> graders in Faribault County in 2016. Use among 9<sup>th</sup> graders has remained relatively steady since 1995, with the highest level (12%) in 2013.

### Prescription Drug Misuse

Past 30-day prescription drug misuse was reported by 2% of 8<sup>th</sup> graders, 9% of 9<sup>th</sup> graders, and 8% of 11<sup>th</sup> graders in Faribault County in 2016. Specifically, in the past year, students in these grades combined reported the misuse of stimulants (1.9%); ADD or ADHD medication (4.8%); pain relievers (6.1%); and tranquilizers (2.9%).

**Figure 2. Faribault County 11<sup>th</sup> Graders Reporting How Much They Typically Drink at One Time (2016 MSS)**



Mental health disorders can increase risk of substance use, and substance use can in turn increase the risk for poor mental health. In addition, numerous risk and protective factors are shared between substance use and mental health outcomes.

To learn more, read the SUMN fact sheet on this topic:

[http://sumn.org/~media/498/MENTAL\\_HEALTH\\_2017\\_SUMN\\_Fact\\_Sheet.pdf](http://sumn.org/~media/498/MENTAL_HEALTH_2017_SUMN_Fact_Sheet.pdf)

## Contributing Factors

Low perception of harm, low perception of parental disapproval, and misperception of peer use all contribute to alcohol, tobacco, and drug use in a community. Although data are presented here as a group, percentages vary by grade. For data by grade level, or for more protective factors, visit [www.sumn.org](http://www.sumn.org).

### Perceived Harm

Faribault County 5<sup>th</sup>, 8<sup>th</sup>, 9<sup>th</sup>, and 11<sup>th</sup> graders were least likely to believe that people put themselves at great or moderate risk of harm, physically and in other ways, by smoking marijuana once or twice per week: 61%. Perceived great or moderate risk of harm for other substances was as follows: consuming five or more alcoholic drinks once or twice per week: 67%; smoking one or more packs of cigarettes per day: 79%; using prescription drug not prescribed for you: 78%.

### Perceived Disapproval

Among Faribault County 5<sup>th</sup>, 8<sup>th</sup>, 9<sup>th</sup>, and 11<sup>th</sup> graders, 87% said their parents would think it's very wrong or wrong for them to smoke cigarettes. Similarly, students felt their parents would feel it's very wrong or wrong for them to have one or two drinks of alcohol nearly every day (87%), to smoke marijuana (89%) or to use prescription drugs not prescribed for them (92%). Faribault County students also believe their friends would disapprove of substance use: most students believe their friends would think it was wrong for them to engage in cigarette use (79%); alcohol use (75%); marijuana use (72%); or prescription drug misuse (85%).

### School Factors

Faribault County 8<sup>th</sup>, 9<sup>th</sup>, and 11<sup>th</sup> graders with better student-teacher relationships are 2.5 times less likely to report past-month marijuana use, and those with greater educational engagement were nearly 3 times less likely. Similar associations are seen with alcohol, tobacco, and prescription drugs.

### Misperception of Peer Use

Faribault County 8<sup>th</sup>, 9<sup>th</sup>, and 11<sup>th</sup> graders who think most students at their school drink monthly or more often are nearly 9 times more likely to report that they themselves drink alcohol monthly or more often.

**Table 2. Perceived vs. Actual Use, Faribault County 2016 Minnesota Student Survey**

	8 <sup>th</sup> Grade	9 <sup>th</sup> Grade	11 <sup>th</sup> Grade
Students reporting they think MOST students in their school drink alcohol once per month or more often	24%	50%	82%
Students reporting they drink alcohol themselves once per month or more often	2%	5%	23%
Students reporting they think MOST students in their school use marijuana once per month or more often	26%	53%	77%
Students reporting they use marijuana themselves once per month or more often	0%	7%	15%

## Consequences

### Alcohol


- The DWI arrest rate for Faribault County residents has remained relatively stable since 2006, averaging 52.6 per 10,000 with 48.8 in 2015. The state rate has decreased since 2007, with an average of 61.8 per 10,000 population since 2006, and 44.7 in 2015 (Minnesota Office of Traffic Safety, DPS)
- In 2015, 8.3% of motor vehicle crashes in Faribault County were alcohol-related. In comparison, 4.9% throughout the state were alcohol-related (*Crash Facts*, DPS)

### Tobacco

- The lung, bronchus and trachea cancer death rate for Faribault County has fluctuated between 5.5 and 11.4 per 10,000 population between 2006 and 2015, averaging 8.4. That's higher than the state average of 4.5 per 10,000 (Minnesota Center for Health Statistics, MDH)

### Drugs

- From 2000 to 2015, there were 3 opioid deaths (non-exclusive) in Faribault County, of 2,273 in Minnesota. Over that same 15-year period, there was 1 heroin death in Faribault County; there were 453 in the state as a whole (Minnesota Center for Health Statistics, MDH)
- In 2015, there were 4 arrests for marijuana in Faribault County, no arrests for opium or cocaine, 4 arrests for synthetic drugs, and 3 arrests for other drugs (*Minnesota Crime Information*, DPS)



**Figure 3. Admissions to Treatment for Opioids as the Primary Substance of Abuse among Faribault County Residents Admitted (Drug and Alcohol Abuse Normative Evaluation System)**

### SUMN.org

SUMN is a one-stop-shop for data, tools, and prevention resources. Visitors can search county, regional, and state level data by topic, by location, and by demographic producing tables, charts, graphs, and maps. View and download data products, such as this fact sheet, and find tips on finding, analyzing, translating, using, and disseminating data.

- Use the 'Data by Topic' search function to access county level data, by grade and gender, for all measures included in this fact sheet.
- Visit the 'Publications' page to find SUMN fact sheets on a variety of risk and protective factors, including educational engagement and teacher-student relationships.

Funded by the Minnesota Department of Human Services Alcohol and Drug Abuse Division

## Contact Us

For questions or more information regarding the State Epidemiological Outcomes Workgroup or SUMN.org, please email: [info@sumn.org](mailto:info@sumn.org)

To contact the Regional Prevention Coordinator for your region, please visit: [www.rpcmn.org](http://www.rpcmn.org)