



POPE COUNTY

SUMN.org FACT SHEET

2017

Substance Abuse Prevention

This fact sheet provides county level data on substance use and related factors to help guide decisions about prevention efforts. These data can be used to justify need for prevention funding, monitor substance-related trends, and plan programs and initiatives by establishing community level prevention priorities.


According to 2016 estimates from the U.S. Census Bureau, Pope County has 11,049 residents; 21% are under the age of 18.

Key Findings

- Compared to state averages, smaller percentages of Pope County students reported past 30-day tobacco use and marijuana use (see Fig. 1).
- Cigarette smoking among 9th graders dropped in the county from 16% in 1998 to 6% in 2016; however, 4% of 9th graders also reported using e-cigarettes in 2016.
- Among Pope County's 8th, 9th, and 11th graders, those with better teacher-student relationships are over 2 times less likely to report past-month prescription drug misuse.
- In 2015, Pope County's alcohol-related motor vehicle crash rate was higher than the state rate, but the arrest rate for driving while intoxicated (DWI) was lower.
- From 2000 to 2015, Pope County recorded 1 heroin overdose death due to heroin, and none due to opioids.

For more information on alcohol, tobacco, and other drug use, consequences, contributing factors, and treatment, as well as mental health data, visit SUMN.org

Figure 1. 8th, 9th, and 11th Grade Past 30-Day Substance Use, 2016 Minnesota Student Survey (MSS)



**Tobacco includes any past month use of cigarettes, cigars, chewing tobacco, electronic cigarettes and/or hookah.

Adult Substance Use

According to the 2015 Minnesota Survey on Adult Substance Use (MNSASU):

- Past-month binge drinking (women having 4+ drinks or men having 5+ drinks in a row on one occasion) was reported by 13% of adults in the West Central region and 14% of Minnesota adults.
- Past-month cigarette smoking was reported by 19% of West Central adults and 16% of Minnesota adults.
- Past-month marijuana use was reported by 3% of adults in the West Central region compared to 5% of adults statewide.

Youth Substance Use

Data on youth substance use are from the Minnesota Student Survey (MSS); Pope County students make up 0.2% of the overall statewide sample.

	5 th Grade	8 th Grade	9 th Grade	11 th Grade
Females	35	32	25	24
Males	40	43	30	38
No Answer	1	1	0	0
Total	76	76	55	62

Past year alcohol and/or drug use was reported by 4.1% of Pope 5th graders. Questions about past 30-day substance use (covered in the following section) are not asked of 5th grade students.

Alcohol

Past 30-day alcohol use was reported by 13% of 8th graders, 13% of 9th graders, and 15% of 11th graders in Pope County in 2016. Use among 9th graders dropped in Pope County from 27% in 1998 to 13% in 2016. Past 30-day binge drinking (defined as having five or more drinks in a row on one occasion) was reported by 5% of 8th graders, 6% of 9th graders, and 7% of 11th graders in 2016.

Tobacco/Nicotine

Of Pope County 8th, 9th and 11th graders, 11% reported any past 30-day tobacco use: 7% reported smoking cigarettes, 3% reported smoking cigars or cigarillos, and 5% reported chewing tobacco use. In addition, 6% reported e-cigarette use, and 1% reported hookah use; questions about these two methods of nicotine use were added to the survey in 2016. Cigarette smoking among 9th graders dropped in Pope County from 16% in 1998 to 6% in 2016.

Marijuana

Past 30-day marijuana use was reported by 7% of 8th graders, 9% of 9th graders, and 8% of 11th graders in Pope County in 2016. Use among 9th graders decreased in Pope County from 10% in 1998 to 9% in 2016.

Prescription Drug Misuse

Past 30-day prescription drug misuse was reported by 4% of 8th graders, 7% of 9th graders, and 7% of 11th graders in Pope County in 2016. Specifically, in the past year, students in these grades combined reported the misuse of stimulants (1.6%); ADD or ADHD medication (1.6%); pain relievers (2.7%); and tranquilizers (2.1%).

Figure 2. Pope County 11th Graders Reporting How Much They Typically Drink at One Time (2016 MSS)



Mental health disorders can increase risk of substance use, and substance use can in turn increase the risk for poor mental health. In addition, numerous risk and protective factors are shared between substance use and mental health outcomes.

To learn more, read the SUMN fact sheet on this topic:

http://sumn.org/~media/498/MENTAL_HEALTH_2017_SUMN_Fact_Sheet.pdf

Contributing Factors

Low perception of harm, low perception of parental disapproval, and misperception of peer use all contribute to alcohol, tobacco, and drug use in a community. Although data are presented here as a group, percentages vary by grade. For data by grade level, or for more protective factors, visit www.sumn.org.

Perceived Harm

Pope County 5th, 8th, 9th, and 11th graders were least likely to believe that people put themselves at great or moderate risk of harm, physically and in other ways, by smoking marijuana once or twice per week: 66%. Perceived great or moderate risk of harm for other substances was as follows: five or more alcoholic drinks once or twice per week: 72%; smoking one or more packs of cigarettes per day: 83%; using prescription drug not prescribed for you: 85%.

Perceived Disapproval

Among Pope County 5th, 8th, 9th, and 11th graders, 93% said their parents would think it's very wrong or wrong for them to smoke cigarettes. Similarly, students felt their parents would feel it's very wrong or wrong for them to have one or two drinks of alcohol nearly every day (88%), to smoke marijuana (90%) or to use prescription drugs not prescribed for them (95%). Pope County students also believe their friends would disapprove of substance use: most students believe their friends would think it was wrong for them to engage in cigarette use (87%); alcohol use (86%); marijuana use (85%); or prescription drug misuse (92%).

School Factors

Pope County 8th, 9th, and 11th graders with better student-teacher relationships are over 2 times less likely to report past-month prescription drug misuse, and those with greater educational engagement were nearly 2 times less likely. Similar associations are seen with alcohol, tobacco, and marijuana.

Misperception of Peer Use

Pope County 8th, 9th, and 11th graders who think most students at their school drink monthly or more often are over 4 times as likely to report that they themselves drink alcohol monthly or more often.

Table 2. Perceived vs. Actual Use, Pope County 2016 Minnesota Student Survey

	8 th Grade	9 th Grade	11 th Grade
Students reporting they think MOST students in their school drink alcohol once per month or more often	32%	50%	63%
Students reporting they drink alcohol themselves once per month or more often	5%	11%	15%
Students reporting they think MOST students in their school use marijuana once per month or more often	24%	33%	60%
Students reporting they use marijuana themselves once per month or more often	3%	9%	5%

Consequences

Alcohol


- The DWI arrest rate for Pope County residents fell steadily between 2006 and 2011, from 60.7 per 10,000 population to 34.8. It has risen slightly since then, to 39.0 in 2015, with an average of 44.4 from 2006 to 2015. The state rate followed a similar pattern but has been slightly higher than Pope's over time, with an average of 57.0 (Minnesota Office of Traffic Safety, DPS)
- In 2015, 11.9% of motor vehicle crashes in Pope County were alcohol-related. In comparison, 4.9% throughout the state were alcohol-related (*Crash Facts*, DPS)

Tobacco

- The lung, bronchus and trachea cancer death rate for Pope County has fluctuated between 9.0 and 3.7 per 10,000 population, from 2006 to 2015, averaging 6.8 per 10,000 population. That's higher than the state average of 4.5 per 10,000 (Minnesota Center for Health Statistics, MDH)

Drugs

- From 2000 to 2015, while there were 2,273 opioid deaths (non-exclusive) in Minnesota, none of those deaths occurred in Pope County. Over that same 15-year period, there was 1 heroin death in Pope County, which occurred in 2014. (Minnesota Center for Health Statistics, MDH)
- In 2015, there were 4 arrests for marijuana in Pope County, 1 arrest for opium or cocaine, 2 for synthetic drugs, and 19 arrests for other drugs, including methamphetamine (*Minnesota Crime Information*, DPS)



Percentage of Opioid Admissions to Minnesota Treatment Facilities for Drug Use by Pope County Residents (The Drug and Alcohol Abuse Normative Evaluation System, Minnesota Department of Human Services)

State Epidemiological Outcomes Workgroup

Minnesota's SEOW has been monitoring substance abuse trends since 2007. The Workgroup helps guide prevention planning at the state level, and provides training and technical assistance to communities. In 2008 the SEOW launched the Substance Use in Minnesota (SUMN) website to ensure quick and easy access to data for prevention professionals.

SUMN.org

SUMN is a one-stop-shop for data, tools, and prevention resources. Visitors can search county, regional, and state level data by topic, by location, and by demographic producing tables, charts, graphs, and maps. View and download data products, such as this fact sheet, and find tips on finding, analyzing, translating, using, and disseminating data.

Contact Us

For questions or more information regarding the State Epidemiological Outcomes Workgroup or SUMN.org, please email: info@sumn.org

To contact the Regional Prevention Coordinator for your region, please visit: www.rpcmn.org

Funded by the Minnesota Department of Human Services Alcohol and Drug Abuse Division